



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2014

---

## **Alien interference: disruption of infochemical networks by invasive insect herbivores**

Desurmont, Gaylord A ; Harvey, Jeff ; van Dam, Nicole M ; Cristescu, Simona M ; Schiestl, Florian P ; Cozzolino, Salvatore ; Anderson, Peter ; Larsson, Mattias C ; Kindlmann, Pavel ; Danner, Holger ; Turlings, Ted C J

**Abstract:** Insect herbivores trigger various biochemical changes in plants, and as a consequence, affect other organisms that are associated with these plants. Such plant-mediated indirect effects often involve herbivore-induced plant volatiles (HIPVs) that can be used as cues for foraging herbivores and their natural enemies, and are also known to affect pollinator attraction. In tightly co-evolved systems, the different trophic levels are expected to display adaptive response to changes in HIPVs caused by native herbivores. But what if a new herbivore invades such a system? Current literature suggests that exotic herbivores have the potential to affect HIPV production, and that plant responses to novel herbivores are likely to depend on phylogenetic relatedness between the invader and the native species. Here we review the different ways exotic herbivores can disrupt chemically mediated interactions between plants and the key users of HIPVs: herbivores, pollinators, and members of the third (i.e. predators and parasitoids) and fourth (i.e. hyperparasitoids) trophic levels. Current theory on insect invasions needs to consider that disruptive effects of invaders on infochemical networks can have a short-term impact on the population dynamics of native insects and plants, as well as exerting potentially negative consequences for the functioning of native ecosystems.

DOI: <https://doi.org/10.1111/pce.12333>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-96076>

Journal Article

Accepted Version

Originally published at:

Desurmont, Gaylord A; Harvey, Jeff; van Dam, Nicole M; Cristescu, Simona M; Schiestl, Florian P; Cozzolino, Salvatore; Anderson, Peter; Larsson, Mattias C; Kindlmann, Pavel; Danner, Holger; Turlings, Ted C J (2014). Alien interference: disruption of infochemical networks by invasive insect herbivores. *Plant, Cell Environment*, 37(8):1854-1865.

DOI: <https://doi.org/10.1111/pce.12333>

# Alien interference: Disruption of infochemical networks by invasive insect herbivores

Gaylord A. Desurmont<sup>1\*</sup>, Jeff Harvey<sup>2</sup>, Nicole M. van Dam<sup>3</sup>, Simona Cristescu<sup>4</sup>, Florian P. Schiestl<sup>5</sup>,  
Salvatore Cozzolino<sup>6</sup>, Peter Anderson<sup>7</sup>, Mattias C. Larsson<sup>7</sup>, Pavel Kindlmann<sup>8</sup>, Holger Danner<sup>3</sup>,  
Ted C.J. Turlings<sup>1</sup>

<sup>1</sup>Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

<sup>2</sup>Terrestrial Ecology, Netherlands Institute of Ecology, Wageningen, The Netherlands

<sup>3</sup>Ecogenomics, Radboud University Nijmegen, Nijmegen, The Netherlands

<sup>4</sup>Institute for Molecules and Materials, Radboud University Nijmegen, Nijmegen, The Netherlands

<sup>5</sup>Institut für Systematische Botanik, University of Zurich, Zurich, Switzerland

<sup>6</sup>Dipartimento di Biologia, Università degli Studi di Napoli Federico II, Napoli, Italy

<sup>7</sup>Plant Protection Biology, Swedish University of Agricultural Sciences, Alnarp, Sweden

<sup>8</sup>Institute for Environmental Studies, Charles University, Prague, Czech Republic

\*Corresponding author:

Gaylord A. Desurmont, Institute of Biology, University of Neuchâtel, Rue Emile-Argand 11, CH-2000  
Neuchâtel, Switzerland, gaylord.desurmont@unine.ch

Running Title: *Invasive insects and infochemicals*

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/pce.12333

## ABSTRACT

Insect herbivores trigger various biochemical changes in plants and, as a consequence, affect other organisms that are associated with these plants. Such plant-mediated indirect effects often involve herbivore-induced plant volatiles (HIPVs) that can be used as cues for foraging herbivores and their natural enemies, and are also known to affect pollinator attraction. In tightly co-evolved systems, the different trophic levels are expected to display adaptive response to changes in HIPVs caused by native herbivores. But what if a new herbivore invades such a system? Current literature suggests that exotic herbivores have the potential to affect HIPV production, and that plant responses to novel herbivores are likely to depend on phylogenetic relatedness between the invader and the native species. Here we review the different ways exotic herbivores can disrupt chemically mediated interactions between plants and the key users of HIPVs: herbivores, pollinators, and members of the third (i.e. predators and parasitoids) and fourth (i.e. hyperparasitoids) trophic levels. Current theory on insect invasions needs to consider that disruptive effects of invaders on infochemical networks can have a short term impact on the population dynamics of native insects and plants, as well as exerting potentially negative consequences for the functioning of native ecosystems.

*Key-words:* invasive species, plant volatiles, multitrophic interactions, plant-insect interactions

47

## 48 INTRODUCTION

49

50 Plant volatiles are of key importance for the foraging behaviour of many insects.

51 Pollinators, for example, use plant-produced volatiles to locate rewarding flowers

52 (Pichersky & Gershenzon, 2002, Raguso, 2008, Schiestl, 2010). In contrast to these

53 mutualists, many herbivores utilize plant volatiles to identify suitable host plants

54 (Bruce, Wadhams & Woodcock, 2005). Of particular interest is the phenomenon that

55 plants under attack by arthropod herbivores show dramatic increases in the production

56 of specific volatiles. Certain herbivores are deterred by these inducible volatiles (De

57 Moraes, Mescher & Tumlinson, 2001, Kessler & Baldwin, 2001), possibly to avoid

58 competition, reduced food quality, or natural enemies, but others are attracted to

59 already infested plants (Heil, 2004, Horiuchi, Arimura, Ozawa, Shimoda, Dicke,

60 Takabayashi & Nishioka, 2003), in certain cases to initiate mass attacks on weakened

61 hosts (Raffa, Phillips & Salom, 1993).

62 Inducible plant volatiles are also of great importance to predators and

63 parasitoids in search of prey or hosts. Therefore, herbivore-induced plant volatiles

64 (HIPVs) are proposed to function as a form of indirect resistance that relies on the third

65 trophic level to reduce herbivore damage (Dicke & Sabelis, 1988, Turlings, Tumlinson &

66 Lewis, 1990). This indirect defensive function is still highly debated (Allison & Hare,

67 2009, Arimura, Matsui & Takabayashi, 2009, Hare, 2011, Heil & Karban, 2010,

68 Holopainen, 2004, Peñuelas & Llusà, 2004), but there is an overall consensus that

69 arthropods at different trophic levels use plant volatiles to locate key resources and that

70 these chemically-mediated interactions may be very specific, with each species having

71 evolved to exploit particular volatile blends. We propose that invading insect herbivores

may disrupt these finely-tuned chemically mediated interactions and that this may have consequences for the reproductive success and eventually the population structure of the interacting native plants and insects.

With increasing globalization, many plants and animals are being accidentally or intentionally introduced into non-native ecosystems, and some of these invaders pose major threats for biodiversity and the functioning of ecological processes in their introduced range (Gandhi & Herms, 2010, Vilà, Espinar, Hejda, Hulme, Jarošík, Maron, Pergl, Schaffner, Sun & Pyšek, 2011). Moreover, recently observed changes in climate that are driven largely by humans are also challenging species to adapt and many are responding by moving polewards. The consequences of these invasions have been studied extensively, but very little attention has been given to many important mechanisms involved, such as plant odour-mediated interactions (Gandhi & Herms, 2010, Kenis, Auger-Rozenberg, Roques, Timms, Péré, Cock, Settele, Augustin & Lopez-Vaamonde, 2009). In the context of a EUROCORE project we recently launched a collaborative effort to study this concept of alien herbivores interfering with infochemical networks (Fig. 1). We particularly focus on quantifying the effects of invasive insect herbivores on the foraging behaviour of native herbivores and plant mutualists, such as pollinators and parasitic wasps. To our knowledge, no study has specifically addressed this issue, but a number of investigations on the interactions among assemblages of native arthropods (Bukovinszky, Poelman, Kamp, Hemerik, Prekatsakis & Dicke, 2012), or between native arthropods and invasive plants (Harvey & Fortuna, 2012) strongly suggest that the disruption of plant–odour mediated interactions can have ecological consequences (Table 1). Within this framework, a key point to address is whether and how the effects of exotic herbivores on HIPVs differ from effects caused by native herbivores. Indeed, native species can also play the role of

97 “info-disrupter” in interactions within native communities. For example, the effect of  
98 *Mamestra brassicae*, a Eurasian moth, on the volatiles emitted by crucifers was shown to  
99 influence the outcome of the interactions between another native herbivore, *Pieris*  
100 *brassicae*, and its parasitoid *Cotesia glomerata* (Bukovinszky *et al.*, 2012). Thus,  
101 demonstrating that invasive species have an impact on infochemical networks may not  
102 be a sufficient conclusion: the ultimate ecological challenge is to determine whether or  
103 not the impact of invaders is greater than or different from the impact naturally caused  
104 by other members of the native fauna, and poses a greater threat to the functioning of  
105 native ecosystems. In this context, a comparative approach would be most appropriate.  
106 Studying single pairs of organisms (one native species/one exotic species) may be a  
107 valuable approach in a specific context, for example predicting the outcome of the  
108 association between a threatening invasive species and a particularly rare or  
109 ecologically important native species. However, to reveal general ecological trends and  
110 to answer the question of the global impact of invasive species on infochemical  
111 networks, a standardized comparison of the effects of several native and exotic species  
112 on chemically mediated interactions will be necessary.

113 Two fundamental questions pertain to the notion of interference of chemically  
114 mediated interactions by invasive insect herbivores: 1) how do plants perceive and  
115 respond to “novel” herbivory? And 2) are insects that rely on infochemical networks  
116 adapted to variations in HIPVs and able to cope with those induced by invasive  
117 herbivores? Here we combine our current knowledge of biochemical mechanisms  
118 involved in the production of plant volatiles and recent invasion biology literature to  
119 unite/bridge both questions. Then, we address the question of adaptation to variation in  
120 HIPVs for each of the trophic levels involved in infochemical networks: herbivores,

121 natural enemies of herbivores (third trophic level) and their own natural enemies (i.e.  
122 hyperparasitoids, fourth trophic level), and pollinators.

123

#### 124 **HOW DO PLANTS PERCEIVE AND RESPOND TO “NOVEL” HERBIVORY?**

125

126 The notion of novel herbivory, hereby defined as feeding by an herbivore that the plant  
127 has not experienced over evolutionary time, is relative. Due to the immense variety of  
128 herbivores attacking plants worldwide and the diversity of feeding strategies they  
129 occupy, the possibility of an invasive insect attacking and damaging native plants in an  
130 entirely novel way is unlikely. Geographically isolated environments, such as islands,  
131 where plants could have existed in absence of some specific guilds of herbivores

132 (Reaser, Meyerson, Cronk, De Poorter, Eldrege, Green, Kairo, Latasi, Mack &  
133 Mauremootoo, 2007), may be an exception to that rule, but in the vast majority of exotic  
134 insects invasions, the invader will become an additional species integrated into native  
135 plant-insect food webs. Still, the scenario of an invasive species attacking native plants  
136 in the exact same way as a native herbivore is equally unlikely, because modes of  
137 feeding vary tremendously from species to species. Preferences for specific plant tissues  
138 or organs, feeding patterns, activity rhythms and circadian cycles, solitary or gregarious  
139 feeding behaviours, chemistry of saliva, and counter-adaptations to plant defences are  
140 among the many parameters that potentiate species-specific variation. In other words,  
141 all exotic herbivores are expected to have some “elements of novelty”, i.e. diverge from  
142 existing native herbivores in some specific traits, but none, or very few, will pose an  
143 entirely new threat to native plants. The true question is to know which herbivore traits  
144 are fundamental to the perception by the plant of herbivory and its response in the form

of a volatile blend emitted upon attack, and how these traits are expected to differ between native and exotic herbivores.

#### *Herbivore cues recognized by plants*

In order to investigate whether or not plant responses to exotic herbivores may differ from responses to native herbivores, we must first understand how plants perceive herbivorous arthropods, a topic that has been the subject of several recent reviews (Hilker & Meiners, 2010, Mithöfer & Boland, 2012, Mumm & Dicke, 2010, Wu & Baldwin, 2010). It is well known that plants typically produce different blends of HIPVs when damaged by insects compared to plants that are merely mechanically wounded, implying the existence of insect-derived compounds (including plant compounds modified by insect enzymes) responsible for initiating specific plant responses. Such elicitors have been discovered in insect oral secretions for several systems (Pare, Farag, Krishnamachari, Zhang, Ryu & Kloepper, 2005, Tumlinson & Engelberth, 2008). Additionally, plants can perceive other types of insect cues, related to touch (Hall, MacGregor, Nijse & Bown, 2004), or associated with oviposition (Reymond, 2013). Plant responses to oviposition-related elicitors range from direct defence [e.g. necrosis, growth of undifferentiated wound tissue (Balbyshev & Lorenzen, 1997, Desurmont & Weston, 2011)] to the production of specific HIPV blends attractive to natural enemies (Fatouros, Bukovinskine’Kiss, Dicke & Hilker, 2007).

While critical for plant responses, biochemical elicitors do not constitute the only facet of herbivory. The type of damage inflicted to the plant is also known to influence the production of plant volatiles. Sucking piercing insects such as aphids induced different responses than chewing insects such as caterpillars (Bidart-Bouzat & Kliebenstein, 2011). Also the location (i.e. organ attacked) and the “physical



characteristics" (e.g. scratching, chewing, or skeletonizing damage for leaf-feeders) of the damage may matter and result in the production of distinct blends of HIPVs (Mithöfer & Boland, 2008, Pierre, Jansen, Hordijk, van Dam, Cortesero & Dugravot, 2011). This matter is complicated by the fact that feeding habits can be plastic and change over the lifetime of an herbivore, as illustrated by *P. brassicae* caterpillars, which are typically leaf feeders but switch to flower-feeding during their development on *Brassica nigra* (Smallegange, Van Loon, Blatt, Harvey, Agerbirk & Dicke, 2007). Finally, plants can respond to the time course and circadian rhythm of herbivory (Arimura, Köpke, Kunert, Volpe, David, Brand, Dabrowska, Maffei & Boland, 2008, Connor, Rott, Samietz & Dorn, 2007, Mithöfer, Wanner & Boland, 2005). These results suggest that the production of specific volatile blends in response to herbivory can be, in certain cases, independent of the presence of elicitors in oral secretions. However, from the plant's perspective, feeding habits may be less reliable of the identity of an attacker than biochemical elicitors due to their potentially variable and dynamic nature.

In summary, we postulate that the volatile blend produced by a plant in response to herbivory can depend on three herbivore-dependent variables: 1) the type of elicitors associated with the herbivore, 2) the type and location of damage inflicted to the plant, and 3) the time course or rhythm of herbivory. Because tremendous variation in these traits exists among potential invasive insect herbivores, making general predictions about the extent to which plant responses to exotic and native herbivores may differ in general is a seemingly impossible task. Practically, in order to predict how a plant will respond to a given exotic species, it is necessary to examine where the novel herbivore is situated within this space of the tripartite herbivory traits, or, in other words, determine its coordinates on a three-dimensional "herbivory map", and compare these coordinates to those of the assemblage of native herbivores associated with the plant.

This trait-matching approach has been used in invasion biology studies, mainly in the context of predicting the native species of herbivores most likely to feed on an exotic plant, and evaluating the risks of non-target effects on native plants following the introduction of a biological control agent of an exotic plant (Pearse, Harris, Karban & Sih, 2013). However, the principal limitation of such approaches is the lack of detailed information available on the traits that matter in the context of a specific interaction. To overcome this obstacle, phylogenetic relatedness has been commonly used as a proxy for similarity in traits or groups of traits between native and exotic species of plants or herbivores (Becerra, 1997, Gilbert, Magarey, Suiter & Webb, 2012).

At a broad scale, related plant species may have similar defensive responses, and exotic herbivore species closely related to native ones may share similar feeding habits.

Interestingly, a phylogenetically controlled comparative study of elicitors of plants defences (FACs) showed that closely related species indeed have more elicitors in common than species that are distantly related (Mori & Yoshinaga, 2011, Yoshinaga, Albourn, Nakanishi, Suckling, Nishida, Tumlinson & Mori, 2010), suggesting evolutionary conservatism of these cues. However, it remains largely unknown how different assemblages of FACs interact and ultimately affect plant response for most herbivores.

In the context of understanding interference effects due to exotic herbivores on infochemical networks, we speculate that the use of phylogenetic tools may be useful to estimate the likelihood of an herbivore or a plant to possess certain traits, but will have limited power to predict the outcome of a novel insect-plant interaction in terms of production of specific plant volatiles after attack, and even less so how these volatiles may interfere in the existing native interactions.

*Conceivable plant responses elicited by exotic herbivores.*

220

221 Alternatively to a predictive, trait-matching approach, a more pragmatic approach may

222 be taken to address the apparent “Pandora’s box” of plant responses to novel

223 herbivores. In the light of our knowledge of plant perception of herbivory, what general

224 types of response can we expect from a plant in terms of divergence from its response to

225 native herbivores? Intuitively, we can think of three possible types of response (Fig. 2):

226 (1) *The null response*. The plant damaged by the exotic herbivore does not release

227 volatiles that differ qualitatively or quantitatively from a mechanically wounded plant.

228 This type of response is most likely to occur if the novel herbivore lacks the chemical

229 elicitors present in the oral secretions of native herbivores. Examination of the elicitors

230 present in a range of lepidopteran herbivores revealed that the most common elicitors

231 (i.e. FACs structurally similar to volicitin) are not present in several species, supporting

232 this notion (Yoshinaga *et al.*, 2010). In a controlled context, comparing the volatile

233 blends emitted in response to plants mechanically wounded with and without oral

234 secretions of the exotic species may be a good way to check whether the plant “reacts” to

235 the novel herbivore. However, given the fact that repeated periodical wounding of a

236 plant can be sufficient to elicit a response different from one-time mechanical damage

237 (Mithöfer *et al.*, 2005), the null response may be hard to define under natural herbivore

238 damage conditions.

239 (2) *The identical response*. The plant damaged by the exotic herbivore emits a blend of

240 volatiles that is undistinguishable from the blend produced in response to a native

241 herbivore. This type of response is more likely to occur if the exotic herbivore is closely

242 related to the native herbivore and overlaps largely in terms of biochemical elicitors and

243 feeding habits.

244 (3) *The novel response*. The plant damaged by the exotic herbivore emits a blend that  
245 differs to some extent from the blends produced in response to its associated native  
246 herbivores. Any type of qualitative or quantitative change in the HIPV blend can  
247 theoretically be considered to have an element of novelty: even subtle differences in  
248 ratios of volatiles have been shown to impact the foraging behaviour of insects (Bruce *et*  
249 *al.*, 2005, Visser & Avé, 1978). The limited number of studies documenting induced  
250 direct defences of plants in response to exotic herbivores shows that different types of  
251 “novel responses” are possible, from reduced defensive response of the plant due to its  
252 evolutionary naiveté (Desurmont, Donoghue, Clement & Agrawal, 2011, Woodard, Ervin  
253 & Marsico, 2012), to an over-sensitive response to a particular herbivore (Radville,  
254 Chaves & Preisser, 2011).

255 The three theoretical types of responses to novel herbivory will have drastically  
256 different implications for insect-plant interactions, depending on the type of interactions  
257 that the plants have with the native insect (antagonistic or mutualistic), and on the  
258 degree to which native insects depend on HIPVs to achieve reproductive success. It is  
259 important to keep in mind that the distinction between an “identical” and a “novel”  
260 response ultimately depends on the level of chemosensitivity of the receiver, and to the  
261 degree of chemical complexity necessary to elicit a behavioural answer. An organism  
262 only able to detect (or only looking for) a reduced number of volatiles compounds may  
263 not distinguish between plants infested by a native and an exotic herbivore if the  
264 differences in HIPVs elicited by both herbivores are minor. On the other hand,  
265 organisms known to change their behaviour in response to subtle chemical changes,  
266 such as certain parasitic wasps, may be able to make this distinction. In other words, the  
267 same HIPVs blend emitted in response to an exotic herbivore may be perceived as an  
268 identical response or a novel response by different receivers.

## IS THERE ADAPTATION TO VARIATION IN HIPVS BY NATIVE INSECTS?

A wide range of insects belonging to different trophic levels and feeding guilds can perceive and make use of plant volatiles induced by herbivory (Dudareva, Negre, Nagegowda & Orlova, 2006, Hare, 2011), but the extent to which HIPVs can affect their reproductive success varies. For insect herbivores and pollinators, HIPVs may carry information regarding the quality of a plant as a resource (Cunningham, Moore, Zalucki & West, 2004, Fitt, 1989), which may result in adaptive attraction or deterrence, but herbivores and pollinators do not depend on HIPVs as long as they can locate suitable undamaged plants/flowers by using constitutively released plant volatiles or other cues, e.g. vision or vibration. On the other hand, specialized natural enemies such as parasitic wasps often rely on HIPVs as key foraging cues in detecting the presence of a host insect on a plant (Godfray, 1994, Quicke, 1997): their ability to detect suitable hosts ultimately determines their fitness. Indeed, parasitoids have been shown to possess a very strong innate propensity to respond to HIPVs (Vet, Lewis & Carde, 1995), as well as a strong capacity to learn novel odours (Turlings, Wäckers, Vet, Lewis & Tumlinson, 1993). Thus, some groups of insects have been under a stronger selective pressure to detect and respond to HIPVs than others, and these groups are at a greater risk to be significantly affected by disturbances in infochemicals caused by invasive herbivores. The other side of that coin is that insects that rely the most on HIPVs are also the most likely to have evolved adaptive responses to variation in HIPVs naturally occurring in their environment. For example, parasitic wasps may benefit from being able to find their hosts on plants growing under a range of environmental conditions, and thus to be adapted to variation in volatiles resulting from plant responses to different abiotic

294 conditions (Chen, Schmelz, Wäckers & Ruberson, 2008, Gouinguené & Turlings, 2002,  
295 Winter & Rostás, 2010). Such adaptations could theoretically make them more “robust”  
296 to potential disturbance effects caused by invaders. In the following sections, we review  
297 the possible ecological consequences of HIPVs interferences for insects belonging to the  
298 different trophic levels involved in infochemical networks, and discuss their potential  
299 for adaptation to these disturbances.

300

## 301 **INSECT HERBIVORES**

302

303 Host plant location and acceptance by insect herbivores is typically performed by  
304 females, and plant chemistry plays an important role in these foraging efforts (Bernays  
305 & Chapman, 1994), not only aboveground but also belowground (Johnson & Nielsen,  
306 2012). Detection of a suitable host-plant may depend on the recognition of specific  
307 chemical compounds (Blight, Pickett, Wadhams & Woodcock, 1995, Nottingham, Hardie,  
308 Dawson, Hick, Pickett, Wadhams & Woodcock, 1991), or of specific ratios of compounds  
309 (Visser & Avé, 1978). Many herbivore species prefer healthy unharmed plants of the  
310 appropriate species, but others prefer to deposit their eggs on plants that have already  
311 been damaged by conspecifics (Desurmont & Weston, 2011, Raffa *et al.*, 1993) or  
312 heterospecific herbivores (Poelman, Broekgaarden, Van Loon & Dicke, 2008, Shiojiri,  
313 Takabayashi, Yano & Takafuji, 2002). The use of HIPVs may be critical in distinguishing  
314 uninfested host plants from infested ones. For example, females of the moth *Heliothis*  
315 *virescens* are repelled by the emission of (Z)-3-hexenyl acetate and other aliphatic esters  
316 of (Z)-3-hexen-1-ol by caterpillar-infested tobacco plants, and prefer to deposit their  
317 eggs on unharmed plants (De Moraes *et al.*, 2001). Both attraction to and avoidance of  
318 infested plants can be adaptive for herbivores (Eigenbrode, Ding, Shiel & Berger, 2002),

319 depending on whether the infested plant becomes more or less susceptible to  
320 subsequent herbivory (Agrawal, 2000, Fordyce, 2003, Karban, Agrawal & Mangel, 1997).  
321 Newly invading insects interfering with these volatile cues may lead to  
322 maladaptive choices by native herbivores, such as attraction to plants with increased  
323 resistance, or the avoidance of perfectly suitable plants. Oviposition “mistakes” (i.e.  
324 phytophagous insects laying eggs on unsuitable plants) have received considerable  
325 attention in the context of plant invasions (Harvey, Bukovinszky & van der Putten, 2010,  
326 Harvey & Fortuna, 2012). Indeed exotic plants stimulating oviposition, but not  
327 supporting offspring development, have been referred to as “ecological traps” for native  
328 herbivores (Keeler & Chew, 2008). Here we advance the idea that exotic insect  
329 herbivores may also lead to suboptimal feeding or oviposition choices and a degradation  
330 of preference/performance relationships for native insects due to HIPVs interferences.  
331 While such effects may naturally occur between pairs of native species within native  
332 communities, we expect invasive exotic herbivores to have a greater disturbance  
333 potential due to the lack of evolutionary history between the invader and native species.  
334 Indeed native species may have evolved adaptive behavioural responses to HIPVs  
335 associated with other native community members, but their responses to “novel” HIPVs  
336 that result from unfamiliar herbivory are unpredictable and thus more likely to be  
337 maladaptive. Ultimately, the realized fitness of native herbivores in presence and  
338 absence of the exotic competitor will determine the strength and direction of the  
339 selection imposed on the foraging behaviour of the native species.  
340 The effects of HIPVs interferences on herbivores may extend beyond the infested  
341 plant and reach neighbouring plants via plant-plant communication. Indeed, it is known  
342 for several systems that plants can perceive and respond to HIPVs emitted by  
343 conspecifics (Arimura, Ozawa, Horiuchi, Nishioka & Takabayashi, 2001) or other plant

species (Karban, 2001), and use these infochemicals to induce or prime their own defences before herbivore attack (Heil & Karban, 2010, Ton, D'Alessandro, Jourdie, Jakab, Karlen, Held, Mauch-Mani & Turlings, 2007). Thus, HIPVs interferences have the potential to not only affect the foraging choices of native herbivores, but also, by changing the defence responses in neighbouring plants, affect herbivore performance and dispersal patterns at the patch level.

Are herbivores well-adapted to HIPVs variations in nature? It is generally assumed that specialized phytophagous insects have developed more finely-tuned interactions with their host plants, and may thus be capable of more complex or subtle choices when evaluating the quality of a resource. For example, the specialist elm leaf beetle *Xanthogaleruca luteola* is attracted to lightly infested elm leaves, but not heavily infested leaves (Meiners, Hacker, Anderson & Hilker, 2005). The same is true for the root-feeding larvae of the maize pest *Diabrotica virgifera virgifera*, which are attracted to already infested root systems, but not when infestation levels are high (Robert, Erb, Hibbard, French, Zwahlen & Turlings, 2012). For the viburnum leaf beetle *Pyrrhalta viburni*, both the magnitude of infestation (quantity of conspecifics' egg masses) and the quality of infestation (number of conspecifics that contributed to the infestation) have an effect on the oviposition preferences of females (Desurmont & Weston, 2010). On the other hand, generalist insect species are also capable of complex foraging decisions, as illustrated by grasshoppers mixing food sources in order to balance their dietary needs (Behmer, 2008, Behmer & Elias, 2000). Overall, there is no clear evidence that olfactory receptor neurons, which are the nerve cells used by insects to detect odours, differ in sensitivity or specificity between specialist and generalist insect herbivores (Bruce *et al.*, 2005). Therefore, diet breadth may not be a reliable indicator of how likely herbivores are to be sensitive or adapted to disturbances in HIPVs. In addition to innate preferences



for certain volatiles, herbivores can also learn from previous experiences: food sources experienced as immature insects have been shown to affect the foraging preferences of adults in several systems (Akhtar & Isman, 2003, Webster, Qvarfordt, Olsson & Glinwood, 2013). Associative learning may thus help overcoming the potential detrimental effects of the presence of exotic herbivores for native species.

A key example of how chemically mediated interactions lead evolutionary trajectories of herbivores in a multitrophic context is given by the two lepidopteran species *Plutella xylostella* and *Pieris rapae*, which coexist on crucifer plants. Plants emit a different odour blend in response to damage by each of these herbivores. The parasitoid *Cotesia vestalis* is attracted by the odours of plants attacked by its host *P. xylostella*, but not by *P. rapae*, which is an unsuitable host and which it therefore does not attack. Females of *P. xylostella* appear to take advantage of that fact by preferring plants attacked by *P. rapae* for oviposition (Shiojiri *et al.*, 2002). These findings illustrate nicely that interference caused by an herbivore (i.e. *Pieris rapae*) on a volatile-mediated interaction (i.e. foraging behaviour of *C. vestalis* on crucifers) can have fitness consequences for a second herbivore (i.e. *Plutella xylostella*), leading to selection for an adaptive response to said interference in the long term.

## INSECT POLLINATORS

Insect-pollinated plants rely on visual and chemical flower signals that help to optimize the attraction of effective pollinators (Fenster, Armbruster, Wilson, Dudash & Thomson, 2004). In many cases, volatiles produced by the flowers play an essential role in attracting pollinators from a distance, and convergent patterns in floral scents have been found among families of flowering plants (Raguso, 2008, Schiestl, 2010). Exotic

herbivores may interfere with this mutualism if the feeding habits of the herbivore (e.g. florivory) directly impact the quality of the floral bouquet, or if plant volatiles emitted after herbivory affect pollinator attraction (Lucas-Barbosa, van Loon & Dicke, 2011). Direct herbivory on flowers has been demonstrated to affect visits by pollinators in several systems (Lohman, Zangerl & Berenbaum, 1996, McCall & Irwin, 2006). On the other hand, the effects of HIPVs emission on pollinator attraction vary: although lower visitation frequency on plants infested by herbivores seems to be the most common scenario (Adler, Karban & Strauss, 2001, Cardel & Koptur, 2010, Danderson & Molano-Flores, 2010, Kessler & Halitschke, 2009, Kessler, Halitschke & Poveda, 2011), higher pollinator attraction has also been documented (Poveda, Steffan-Dewenter, Scheu & Tschardt, 2003). In wild radishes, uninfested plants were more visited by syrphid flies. This preference could not be linked to morphological features of flowers on infested plants (Lehtilä & Strauss, 1997), suggesting that avoidance of infested flowers is based on odorous cues. However, in most cases it is unclear whether the effect of HIPVs on pollination is due to a herbivore-induced volatile in the flower (Kessler & Halitschke, 2009) or to the reduction in the amounts of floral scents released by the plant upon herbivory (Kessler, Diezel & Baldwin, 2010). An interesting aspect is the possible compensation of herbivore-mediated reduction of floral attractiveness. A recent study with *Pieris brassicae* feeding on *Brassica rapa* suggests that plants compensate reduced attractiveness due to lower scent emission with the production of more flowers during early flowering (Schiestl et al in press). This compensatory plant response was not evident when the invasive *Spodoptera littoralis* was feeding on the plant, suggesting stronger fitness impacts of this herbivore-mediated through floral attractiveness to pollinators (Schiestl et al in review). On the long term, a trade-off between defence and reproduction can occur if herbivore-induced volatiles decrease pollination quality, or if

419 floral odours interfere with the efficiency of volatile-mediated defences (i.e. attraction of  
420 herbivore antagonists). The outcome of these interactions for plant fitness will  
421 determine the direction of the investment toward reproduction or defence in presence  
422 of herbivores (Bartkowska & Johnston, 2012, Schiestl & Johnson, 2013).

423 What can we expect from pollinators in terms of adaptation to HIPV variation? As  
424 stated in the introduction, HIPVs may carry information about the quality of a resource  
425 to pollinators, such as the quality and quantity of the nectar reward (Kessler &  
426 Halitschke, 2007), but pollinators will rarely depend solely on the presence of HIPVs to  
427 find flowers. This consideration aside, it is well known that the main groups of generalist  
428 pollinators (honey bees, bumblebees) are remarkable learners (Riffell, 2011), using  
429 experience-based learning of olfactory and visual cues to exploit the most profitable  
430 nectar and pollen sources. They may thus be able to learn quickly whether a plant  
431 infested by an exotic species provides higher or lower quality reward compared to  
432 uninfested plants, and adjust their foraging preferences in accordance (Wright &  
433 Schiestl, 2009).

434 It is hard to predict which general type of plant response to novel herbivory (null,  
435 identical, or novel; Fig 2) is the most likely to have a significant impact on plant-  
436 pollinator interactions. We can only state that novel responses are the most likely to  
437 affect plant-herbivore-pollinator interactions in the invaded environment.

438

#### 439 **INSECTS FROM THE THIRD AND FOURTH TROPHIC LEVELS**

440

441 As the third trophic level we consider the natural enemies of arthropods herbivores.  
442 They can be separated in two main categories: 1) predators, which feed upon a certain  
443 number of prey items during their lifetime, and, 2) parasitic wasps, flies, and a few

species from other orders, also referred to as parasitoids, whose larvae develop on or within eggs, larvae, or adults of insect herbivores, eventually killing their host. Insects belonging to the fourth trophic level are natural enemies of predators and parasitoids.

The best-known insects belonging to the fourth trophic level are hyperparasitoids: parasitic wasps that attack the larvae or pupae of primary parasitoids. In general, parasitoids and hyperparasitoids are much more specialized than predators: they can only parasitize a limited number of host species, less than the number of prey species predators can attack. There are of course exceptions to that rule, and generalist parasitoids and specialized predators are not uncommon (Hassell & May, 1986, Quicke, 1997).

In the search for prey or hosts, predators and parasitoids utilize cues that are directly associated with the presence of hosts (Colazza, Cusumano, Giudice & Peri, 2014) such as chemical traces left by insects' tarsi, frass, honeydew, or silk. These cues are thought to be predominantly used for short-range foraging. To detect prey-infested plants from a distance, natural enemies often rely on plant volatiles released locally and systemically after herbivory (Vet & Dicke, 1992). How reliable are HIPVs as indicators of the presence of specific herbivores on a plant? Reports on the subject (Clavijo McCormick, Unsicker & Gershenson, 2012, Dicke, 1999) indicate considerable variation among systems. On one end of the spectrum, some plants release very similar blends of volatiles in response to attacks by insects from different feeding guilds (Kessler & Baldwin, 2001). On the other end of the spectrum, several plants have been found to release distinct blends of volatiles in response to subtle changes in herbivore attack, such as: developmental stage of the herbivore (Yoneya, Kugimiya & Takabayashi, 2009), density of herbivores (Girling, Stewart-Jones, Dherbecourt, Staley, Wright & Poppy, 2011, Horiuchi, Arimura, Ozawa, Shimoda, Takabayashi & Nishioka, 2003), and whether

herbivores have been parasitized or not (Poelman, Zheng, Zhang, Heemskerk, Cortesero & Dicke, 2011). Such specificity in volatile emission provides reliable information, which can be exploited by parasitoids to optimize their foraging decisions (Poelman *et al.*, 2011). In parallel, hyperparasitoids have recently been found to also use HIPVs to locate their parasitoid hosts. They are able to discriminate not only between plants bearing healthy herbivores and plants bearing parasitized herbivores, but also between plants with herbivores parasitized by their preferred host and plants with herbivores parasitized by a less optimal host (Poelman, Bruinsma, Zhu, Weldegergis, Boursault, Jongema, van Loon, Vet, Harvey & Dicke, 2012). These fascinating findings demonstrate that specific HIPVs can be exploited not only by natural enemies of insect herbivores, but also by the enemies of these enemies, and illustrate the complexity of HIPV-mediated multitrophic interactions.

In this context, volatiles emitted in response to exotic herbivores can interfere with the foraging behaviour of native insects from the third and fourth trophic levels in two main ways: 1) direct attraction to the plant damaged by the exotic herbivore only, and 2) increased or decreased attraction to plants infested by both the exotic herbivore and the native host or prey. The outcome of these two types of interferences on the fitness of predators/parasitoids/hyperparasitoids will also depend on whether the exotic herbivore can be successfully attacked or not. Exotic herbivores that cannot be attacked (=non hosts) constitute a 'dead end' for native natural enemies, and the attraction to plants infested by these non-hosts may be costly in time and energy. Such attraction to non-host infested plants has been documented for native parasitoids (Snoeren, Mumm, Poelman, Yang, Pichersky & Dicke, 2010). On the other hand, exotic herbivores that can be attacked may not negatively affect natural enemies through volatile attraction, but may allow native herbivores to escape predation/parasitism. The

494 consequences of multiple herbivory (here, plants doubly infested with the native and  
495 the exotic herbivore) for parasitoids are much harder to predict and will be highly  
496 system-specific. This topic has been the subject of a recent review (de Rijk, Dicke &  
497 Poelman, 2013).

498 A recent study on the attraction of the predatory mite *Phytoseiulus macropilis*  
499 (Banks) illustrates how an uncommon herbivore can interfere with a chemically-  
500 mediated tritrophic interaction (Fadini, Venzon, Oliveira, Pallini & Vilela, 2010). It was  
501 found that the predator was strongly attracted to plants with the spider mite  
502 *Tetranychus urticae*, but avoided the odour of plants infested with the red mite  
503 *Oligonychus ilicis*. The former is a common, suitable prey for *P. macropilis*, the latter is  
504 not: *O. ilicis* is very rarely found on strawberry and usually does not occur in association  
505 with the predator. Doubly infested plants, carrying both herbivores, were not attractive  
506 to the predator, which supports the notion that interference in the chemical signalling  
507 by the unfamiliar herbivore may occur in this system (Fadini *et al.*, 2010).

508 Specific natural enemies such as the majority of parasitoids are short-lived  
509 organisms that often have a limited temporal and developmental window to find their  
510 hosts in an excessively complex olfactory landscape. Therefore, it can be expected that  
511 they have adapted, not only to the odorous cues betraying the presence of their host, but  
512 also to the potential disturbances in HIPVs that may naturally occur in the native  
513 environment. These adaptations can take two main forms: 1) A complex array of innate  
514 preferences for diverse blends of HIPVs reflecting adaptive choices toward reliable cues  
515 (Clavijo McCormick *et al.*, 2012), and 2) plastic preferences relying on associative  
516 learning to optimize foraging success (Vet & Dicke, 1992). Both of these adaptive  
517 strategies have been reported in predators (Glinwood, Ahmed, Qvarfordt & Ninkovic,  
518 2011) and parasitoids (Geervliet, Vreugdenhil, Dicke & Vet, 1998).

519 Finally, in terms of general plant response to novel herbivory (null, identical, and  
520 novel) and realized impact on native organisms, we can make predictions for insects  
521 belonging to the third and fourth trophic levels based on the notions developed through  
522 this section. First of all, null plant responses are less likely to have an effect, as plant  
523 responses to mechanical damage are usually less intense and less specific compared to  
524 plants damaged by herbivores (Dudareva *et al.*, 2006, Wu & Baldwin, 2010), and  
525 therefore likely to be less attractive to natural enemies. On the other hand, similarity in  
526 response between plants infested by the exotic and the native herbivore may have  
527 critical consequences for natural enemies if the exotic herbivore is a non-host. Indeed  
528 identical responses would make plants infested by hosts and non-hosts  
529 undistinguishable from a distance, with negative consequences for the foraging  
530 efficiency of natural enemies potentially leading to local extinctions due to insufficient  
531 parasitism rates (Vos, Berrocal, Karamaouna, Hemerik & Vet, 2001). Alternatively, if the  
532 exotic herbivore can be attacked by natural enemies, then the presence of the exotic  
533 herbivore on the plant may not negatively affect natural enemies but may still have  
534 consequences for native predator-prey dynamics: on the one hand, native herbivores  
535 may partially escape predation/parasitism if the exotic herbivore becomes a favoured  
536 prey/host item. On the other hand, natural enemy densities may increase due to the  
537 presence of the exotic herbivore as an alternative food, eventually negatively affecting  
538 populations of native herbivores. Finally, the impact of novel responses cannot be  
539 predicted and will certainly vary greatly among systems. The outcome will depend on  
540 the compounds or ratios of compounds that constitute the elements of novelty of the  
541 “exotic blend”. In such cases of information disruption, the consequences for the natural  
542 enemies will strongly depend on their behavioural plasticity and, on the long-term, on  
543 their potential for rapid evolutionary adaptations.

## CONCLUDING REMARKS AND RESEARCH DIRECTIONS

Predicting insect-plant interactions in a rapidly changing world is a challenge that conservation ecologists are facing worldwide (Leimu, Muola, Laukkanen, Kalske, Prill & Mutikainen, 2012, Lüring & Scheffer, 2007, Pearse *et al.*, 2013, White, Wilson & Clarke, 2006). Thus far, the study of invasive species on community-level interactions has been largely confined to the effects of exotic plants on native plant assemblages (Callaway & Aschehoug, 2000, Keane & Crawley, 2002, Sakai, Allendorf, Holt, Lodge, Molofsky, With, Baughman, Cabin, Cohen & Ellstrand, 2001). The introduction of exotic herbivores or natural enemies can also seriously disrupt the functioning of native communities, with effects rippling through the food chain (Fortuna, Vet & Harvey, 2012, Kenis *et al.*, 2009, Pearson & Callaway, 2005). Although these direct and indirect effects of invasive species have been documented, we are not aware of any studies that have specifically addressed the effects of invading insect herbivores on the chemically mediated interactions among plants and native insects. In effect, the presence of an 'interloper' has the potential to diffuse or sever communication channels involving native herbivores, natural enemies, and pollinators, generating spatial asymmetries between different trophic levels that may destabilize tightly linked trophic interactions (Table 1).

Moreover, these effects on infochemical networks are likely to depend on the magnitude of the invasion: if the exotic herbivore becomes more abundant than natives, owing to release from its own specialized natural enemies, then it may exert a disproportionate effect on the native community by inflicting more damage on plants than the native herbivores. In this scenario, volatiles released from plants damaged by the invader may dominate the olfactory landscape, amplifying any interference effect



569 caused by these HIPVs. In closely evolved systems, it can be expected that foraging  
570 insects have adapted their responses to plant-emitted signals to optimize resource  
571 location, and may thus have evolved some adaptive responses to volatile interferences,  
572 but this question has not yet been considered in emerging theory of invasion biology.

573 Regarding the susceptibility of food webs to infochemical disturbances, it is  
574 important to keep in mind that, due to domestication, crop plants may have lost some of  
575 their ability to emit odorous distress signals in response to herbivory, as has been found  
576 for modern varieties of maize that no longer emit root signals that attract  
577 entomopathogenic nematodes (Degenhardt, Hiltpold, Köllner, Frey, Gierl, Gershenzon,  
578 Hibbard, Ellersieck & Turlings, 2009, Rasmann, Köllner, Degenhardt, Hiltpold, Toepfer,  
579 Kuhlmann, Gershenzon & Turlings, 2005) or no longer respond to freshly deposited  
580 moth eggs that may enhance the plants responses to subsequent caterpillar feeding  
581 (Tamiru, Bruce, Woodcock, Caulfield, Midega, Ogol, Mayon, Birkett, Pickett & Khan,  
582 2011). This realization has motivated researchers to restore (Degenhardt *et al.*, 2009),  
583 genetically alter (Kappers, Aharoni, Van Herpen, Luckerhoff, Dicke & Bouwmeester,  
584 2005) or artificially enhance (Christensen, Nemchenko, Borrego, Murray, Sobhy, Bosak,  
585 DeBlasio, Erb, Robert & Vaughn, 2013) the volatile emissions in crop plants. Thus,  
586 agroecosystems often represent simplified olfactory landscapes compared to natural  
587 environments, which may make them even more susceptible to interferences caused by  
588 exotic herbivores.

589 Practically, experimental efforts should be made to understand the likelihood and  
590 magnitude of disruption of infochemical networks by invasive insect herbivores. In the  
591 context of insect invasions, the ecological effects of exotic HIPVs can only be tested by  
592 combining careful examination of plant responses to insect invaders at the biochemical  
593 level and manipulative experiments with native and exotic insects at the community

level. In the end, measurements of fitness (or fitness correlates) of native insect species in presence and in absence of plant volatiles induced by the invader will be the most valuable indicator of the predicted impact of an invasive species as an “info-disrupter” on a given native environment.

Two challenges need to be considered by researchers investigating this topic. Firstly, because the effects of exotic species on the transfer of chemical information are likely to escalate through trophic levels via cascading effects (Harvey *et al.*, 2010), there is a wide variety of interactions involving native organisms that could potentially be impacted by a single invader, making the exhaustive examination of each of them extremely difficult. The challenge lies in determining which interactions should be examined with priority; these decisions should be based on the knowledge of the role and ecological functions of the native species involved in the system studied (Bukovinszky, van Veen, Jongema & Dicke, 2008). Secondly, measurements of fitness correlates of native organisms under controlled laboratory conditions, despite their usefulness, may have limited value compared to results obtained under field conditions. However, working with exotic species under natural conditions can only be done once the exotic organism has started invading the native environment, diminishing the value of risk assessments and impact predictions. To overcome this obstacle, researchers may take advantage of newly invaded environments to conduct field research in a still “naïve” ecosystem, and extrapolate results to areas at risk of being invaded in the future. This approach may be particularly valuable in the case of range-expanding species, owing to global changes such as global warming (Sorte, Williams & Zerebecki, 2010, Walther, Roques, Hulme, Sykes, Pyšek, Kühn, Zobel, Bacher, Botta-Dukát & Bugmann, 2009).

## **Acknowledgements**

The main authors are grateful to the Swiss National Science Foundation, which finances their contribution in the context of the EUROCORES Programme EuroVOL (project InvaVOL) of the European Science Foundation. Thomas Degen ([www.thomas-degen.ch](http://www.thomas-degen.ch)) provided Figure 1.

## References

- Adler L.S., Karban R. & Strauss S.Y. (2001) Direct and indirect effects of alkaloids on plant fitness via herbivory and pollination. *Ecology*, **82**, 2032-2044.
- Agrawal A.A. (2000) Specificity of induced resistance in wild radish: causes and consequences for two specialist and two generalist caterpillars. *Oikos*, **89**, 493-500.
- Akhtar Y. & Isman M. (2003) Larval exposure to oviposition deterrents alters subsequent oviposition behavior in generalist, *Trichoplusia ni* and Specialist, *Plutella xylostella* Moths. *Journal of Chemical Ecology*, **29**, 1853-1870.
- Allison J.D. & Hare D.J. (2009) Learned and naive natural enemy responses and the interpretation of volatile organic compounds as cues or signals. *New Phytologist*, **184**, 768-782.
- Arimura G.-i., Köpke S., Kunert M., Volpe V., David A., Brand P., Dabrowska P., Maffei M.E. & Boland W. (2008) Effects of feeding *Spodoptera littoralis* on lima bean leaves: IV. Diurnal and nocturnal damage differentially initiate plant volatile emission. *Plant Physiology*, **146**, 965-973.
- Arimura G.-i., Matsui K. & Takabayashi J. (2009) Chemical and molecular ecology of herbivore-induced plant volatiles: proximate factors and their ultimate functions. *Plant and Cell Physiology*, **50**, 911-923.

643 Arimura G.-i., Ozawa R., Horiuchi J.-i., Nishioka T. & Takabayashi J. (2001) Plant–plant  
 644 interactions mediated by volatiles emitted from plants infested by spider mites.  
 645 *Biochemical Systematics and Ecology*, **29**, 1049-1061.

646 Balbyshev N.F. & Lorenzen J.H. (1997) Hypersensitivity and egg drop: a novel  
 647 mechanism of host plant resistance to Colorado potato beetle (Coleoptera:  
 648 Chrysomelidae). *Journal of Economic Entomology*, **90**, 652-657.

649 Bartkowska M.P. & Johnston M.O. (2012) Pollinators cause stronger selection than  
 650 herbivores on floral traits in *Lobelia cardinalis* (Lobeliaceae). *New Phytologist*,  
 651 **193**, 1039-1048.

652 Becerra J.X. (1997) Insects on plants: macroevolutionary chemical trends in host use.  
 653 *Science*, **276**, 253-256.

654 Behmer S.T. (2008) Insect herbivore nutrient regulation. *Annual Review of Entomology*,  
 655 **54**, 165.

656 Behmer S.T. & Elias D.O. (2000) Sterol metabolic constraints as a factor contributing to  
 657 the maintenance of diet mixing in grasshoppers (Orthoptera: Acrididae).  
 658 *Physiological and Biochemical Zoology*, **73**, 219-230.

659 Bernays E.A. & Chapman R.F. (1994) *Host-plant selection by phytophagous insects*.  
 660 Springer.

661 Bidart-Bouzat M.G. & Kliebenstein D. (2011) An ecological genomic approach  
 662 challenging the paradigm of differential plant responses to specialist versus  
 663 generalist insect herbivores. *Oecologia*, **167**, 677-689.

664 Blight M., Pickett J., Wadhams L. & Woodcock C. (1995) Antennal perception of oilseed  
 665 rape, *Brassica napus* (Brassicaceae), volatiles by the cabbage seed weevil  
 666 *Ceutorhynchus assimilis* (Coleoptera, Curculionidae). *Journal of Chemical Ecology*,  
 667 **21**, 1649-1664.

668 Bruce T.J., Wadhams L.J. & Woodcock C.M. (2005) Insect host location: a volatile  
 669 situation. *Trends in Plant Science*, **10**, 269-274.

670 Bukovinszky T., Poelman E.H., Kamp A., Hemerik L., Prekatsakis G. & Dicke M. (2012)  
 671 Plants under multiple herbivory: consequences for parasitoid search behaviour  
 672 and foraging efficiency. *Animal Behaviour*, **83**, 501-509.

673 Bukovinszky T., van Veen F.F., Jongema Y. & Dicke M. (2008) Direct and indirect effects  
 674 of resource quality on food web structure. *Science*, **319**, 804-807.

675 Callaway R.M. & Aschehoug E.T. (2000) Invasive plants versus their new and old  
 676 neighbors: a mechanism for exotic invasion. *Science*, **290**, 521-523.

677 Cardel Y.J. & Koptur S. (2010) Effects of florivory on the pollination of flowers: an  
 678 experimental field study with a perennial plant. *International Journal of Plant*  
 679 *Sciences*, **171**, 283-292.

680 Chen Y., Schmelz E.A., Wäckers F. & Ruberson J.R. (2008) Cotton plant, *Gossypium*  
 681 *hirsutum* L., defense in response to nitrogen fertilization. *Journal of Chemical*  
 682 *Ecology*, **34**, 1553-1564.

683 Christensen S.A., Nemchenko A., Borrego E., Murray I., Sobhy I.S., Bosak L., DeBlasio S.,  
 684 Erb M., Robert C.A. & Vaughn K.A. (2013) The maize lipoxygenase, ZmLOX10,  
 685 mediates green leaf volatile, jasmonate and herbivore-induced plant volatile  
 686 production for defense against insect attack. *The Plant Journal*.

687 Clavijo McCormick A., Unsicker S.B. & Gershenzon J. (2012) The specificity of herbivore-  
 688 induced plant volatiles in attracting herbivore enemies. *Trends in Plant Science*,  
 689 **17**, 303-310.

690 Colazza S., Cusumano A., Giudice D.L. & Peri E. (2014) Chemo-orientation responses in  
 691 hymenopteran parasitoids induced by substrate-borne semiochemicals.  
 692 *BioControl*, **59**, 1-17.

693 Connor E.C., Rott A.S., Samietz J. & Dorn S. (2007) The role of the plant in attracting  
 694 parasitoids: response to progressive mechanical wounding. *Entomologia*  
 695 *Experimentalis et Applicata*, **125**, 145-155.

696 Cunningham J.P., Moore C.J., Zalucki M.P. & West S.A. (2004) Learning, odour preference  
 697 and flower foraging in moths. *Journal of Experimental Biology*, **207**, 87-94.

698 Danderson C.A. & Molano-Flores B. (2010) Effects of herbivory and inflorescence size on  
 699 insect visitation to *Eryngium yuccifolium* (Apiaceae) a prairie plant. *The American*  
 700 *Midland Naturalist*, **163**, 234-246.

701 De Moraes C.M., Mescher M.C. & Tumlinson J.H. (2001) Caterpillar-induced nocturnal  
 702 plant volatiles repel conspecific females. *Nature*, **410**, 577-580.

703 de Rijk M., Dicke M. & Poelman E.H. (2013) Foraging behaviour by parasitoids in  
 704 multiherbivore communities. *Animal Behaviour*, **85**, 1517-1528.

705 Degenhardt J., Hiltbold I., Köllner T.G., Frey M., Gierl A., Gershenzon J., Hibbard B.E.,  
 706 Ellersieck M.R. & Turlings T.C. (2009) Restoring a maize root signal that attracts  
 707 insect-killing nematodes to control a major pest. *Proceedings of the National*  
 708 *Academy of Sciences*, **106**, 13213-13218.

709 Desurmont G.A., Donoghue M.J., Clement W.L. & Agrawal A.A. (2011) Evolutionary  
 710 history predicts plant defense against an invasive pest. *Proceedings of the*  
 711 *National Academy of Sciences*, **108**, 7070-7074.

712 Desurmont G.A. & Weston P.A. (2010) Stimuli associated with viburnum leaf beetle  
 713 (*Pyrrhalta viburni*) aggregative oviposition behavior. *Entomologia Experimentalis*  
 714 *et Applicata*, **135**, 245-251.

715 Desurmont G.A. & Weston P.A. (2011) Aggregative oviposition of a phytophagous beetle  
 716 overcomes egg-crushing plant defences. *Ecological Entomology*, **36**, 335-343.

717 Dicke M. (1999) Are herbivore-induced plant volatiles reliable indicators of herbivore  
 718 identity to foraging carnivorous arthropods? In: *Proceedings of the 10th*  
 719 *International Symposium on Insect-Plant Relationships* (eds S. Simpson, A.J.  
 720 Mordue, & J. Hardie), pp. 131-142. Springer Netherlands.

721 Dicke M. & Sabelis M.W. (1988) How Plants Obtain Predatory Mites as Bodyguards.  
 722 *Netherlands Journal of Zoology*, **38**, 148-165.

723 Dudareva N., Negre F., Nagegowda D.A. & Orlova I. (2006) Plant volatiles: recent  
 724 advances and future perspectives. *Critical Reviews in Plant Sciences*, **25**, 417-440.

725 Eigenbrode S.D., Ding H., Shiel P. & Berger P.H. (2002) Volatiles from potato plants  
 726 infected with potato leafroll virus attract and arrest the virus vector, *Myzus*  
 727 *persicae* (Homoptera: Aphididae). *Proceedings of the Royal Society of London.*  
 728 *Series B: Biological Sciences*, **269**, 455-460.

729 Fadini M.A., Venzon M., Oliveira H., Pallini A. & Vilela E.F. (2010) Response of the  
 730 predatory mite *Phytoseiulus macropilis* (Banks) to volatiles produced by  
 731 strawberry plants in response to attack by Tetranychid mites (Acari:  
 732 Phytoseiidae: Tetranychidae). *Neotropical Entomology*, **39**, 248-252.

733 Fatouros N.E., Bukovinskine'Kiss G., Dicke M. & Hilker M. (2007) The response  
 734 specificity of *Trichogramma* egg parasitoids towards infochemicals during host  
 735 location. *Journal of Insect Behavior*, **20**, 53-65.

736 Fenster C.B., Armbruster W.S., Wilson P., Dudash M.R. & Thomson J.D. (2004) Pollination  
 737 syndromes and floral specialization. *Annual Review of Ecology, Evolution, and*  
 738 *Systematics*, 375-403.

739 Fitt G.P. (1989) The ecology of *Heliothis* species in relation to agroecosystems. *Annual*  
 740 *Review of Entomology*, **34**, 17-53.

741 Fordyce J. (2003) Aggregative feeding of pipevine swallowtail larvae enhances hostplant  
 742 suitability. *Oecologia*, **135**, 250-257.

743 Fortuna T.M., Vet L.E. & Harvey J.A. (2012) Effects of an invasive plant on the  
 744 performance of two parasitoids with different host exploitation strategies.  
 745 *Biological Control*, **62**, 213-220.

746 Gandhi K.J. & Herms D.A. (2010) Direct and indirect effects of alien insect herbivores on  
 747 ecological processes and interactions in forests of eastern North America.  
 748 *Biological Invasions*, **12**, 389-405.

749 Geervliet J.B., Vreugdenhil A.I., Dicke M. & Vet L.E. (1998) Learning to discriminate  
 750 between infochemicals from different plant-host complexes by the parasitoids  
 751 *Cotesia glomerata* and *C. rubecula*. *Entomologia Experimentalis et Applicata*, **86**,  
 752 241-252.

753 Gilbert G.S., Magarey R., Suiter K. & Webb C.O. (2012) Evolutionary tools for  
 754 phytosanitary risk analysis: phylogenetic signal as a predictor of host range of  
 755 plant pests and pathogens. *Evolutionary Applications*, **5**, 869-878.

756 Girling R.D., Stewart-Jones A., Dherbecourt J., Staley J.T., Wright D.J. & Poppy G.M. (2011)  
 757 Parasitoids select plants more heavily infested with their caterpillar hosts: a new  
 758 approach to aid interpretation of plant headspace volatiles. *Proceedings of the*  
 759 *Royal Society B: Biological Sciences*, **278**, 2646-2653.

760 Glinwood R., Ahmed E., Qvarfordt E. & Ninkovic V. (2011) Olfactory learning of plant  
 761 genotypes by a polyphagous insect predator. *Oecologia*, **166**, 637-647.

762 Godfray H.C.J. (1994) *Parasitoids: behavioral and evolutionary ecology*. Princeton  
 763 University Press.

764 Gouinguéné S.P. & Turlings T.C. (2002) The effects of abiotic factors on induced volatile  
 765 emissions in corn plants. *Plant Physiology*, **129**, 1296-1307.



766 Hall D.E., MacGregor K.B., Nijse J. & Bown A.W. (2004) Footsteps from insect larvae  
 767 damage leaf surfaces and initiate rapid responses. *European Journal of Plant*  
 768 *Pathology*, **110**, 441-447.

769 Hare J.D. (2011) Ecological role of volatiles produced by plants in response to damage by  
 770 herbivorous insects. *Annual Review of Entomology*, **56**, 161-180.

771 Harvey J.A., Bukovinszky T. & van der Putten W.H. (2010) Interactions between invasive  
 772 plants and insect herbivores: A plea for a multitrophic perspective. *Biological*  
 773 *Conservation*, **143**, 2251-2259.

774 Harvey J.A. & Fortuna T.M. (2012) Chemical and structural effects of invasive plants on  
 775 herbivore–parasitoid/predator interactions in native communities. *Entomologia*  
 776 *Experimentalis et Applicata*, **144**, 14-26.

777 Hassell M. & May R. (1986) Generalist and specialist natural enemies in insect predator-  
 778 prey interactions. *The Journal of Animal Ecology*, 923-940.

779 Heil M. (2004) Direct defense or ecological costs: responses of herbivorous beetles to  
 780 volatiles released by wild lima bean (*Phaseolus lunatus*). *Journal of Chemical*  
 781 *Ecology*, **30**, 1289-1295.

782 Heil M. & Karban R. (2010) Explaining evolution of plant communication by airborne  
 783 signals. *Trends in Ecology & Evolution*, **25**, 137-144.

784 Hilker M. & Meiners T. (2010) How do plants “notice” attack by herbivorous arthropods?  
 785 *Biological Reviews*, **85**, 267-280.

786 Holopainen J.K. (2004) Multiple functions of inducible plant volatiles. *Trends in Plant*  
 787 *Science*, **9**, 529-533.

788 Horiuchi J.-I., Arimura G.-I., Ozawa R., Shimoda T., Dicke M., Takabayashi J. & Nishioka T.  
 789 (2003) Lima bean leaves exposed to herbivore-induced conspecific plant

790 volatiles attract herbivores in addition to carnivores. *Applied Entomology and*  
791 *Zoology*, **38**, 365-368.

792 Horiuchi J.-I., Arimura G.-I., Ozawa R., Shimoda T., Takabayashi J. & Nishioka T. (2003) A  
793 comparison of the responses of *Tetranychus urticae* (Acari: Tetranychidae) and  
794 *Phytoseiulus persimilis* (Acari: Phytoseiidae) to volatiles emitted from lima bean  
795 leaves with different levels of damage made by *T. urticae* or *Spodoptera exigua*  
796 (Lepidoptera: Noctuidae). *Applied Entomology and Zoology*, **38**, 109-116.

797 Johnson S.N. & Nielsen U.N. (2012) Foraging in the dark—chemically mediated host plant  
798 location by belowground insect herbivores. *Journal of Chemical Ecology*, **38**, 604-  
799 614.

800 Kappers I.F., Aharoni A., Van Herpen T.W., Luckerhoff L.L., Dicke M. & Bouwmeester H.J.  
801 (2005) Genetic engineering of terpenoid metabolism attracts bodyguards to  
802 *Arabidopsis*. *Science Signaling*, **309**, 2070.

803 Karban R. (2001) Communication between sagebrush and wild tobacco in the field.  
804 *Biochemical Systematics and Ecology*, **29**, 995-1005.

805 Karban R., Agrawal A.A. & Mangel M. (1997) The benefits of induced defenses against  
806 herbivores. *Ecology*, **78**, 1351-1355.

807 Keane R.M. & Crawley M.J. (2002) Exotic plant invasions and the enemy release  
808 hypothesis. *Trends in Ecology & Evolution*, **17**, 164-170.

809 Keeler M.S. & Chew F.S. (2008) Escaping an evolutionary trap: preference and  
810 performance of a native insect on an exotic invasive host. *Oecologia*, **156**, 559-  
811 568.

812 Kenis M., Auger-Rozenberg M.-A., Roques A., Timms L., Péré C., Cock M.J., Settele J.,  
813 Augustin S. & Lopez-Vaamonde C. (2009) Ecological effects of invasive alien

814 insects. In: *Ecological Impacts of Non-Native Invertebrates and Fungi on*  
815 *Terrestrial Ecosystems*, pp. 21-45. Springer.

816 Kessler A. & Baldwin I.T. (2001) Defensive function of herbivore-induced plant volatile  
817 emissions in nature. *Science*, **291**, 2141-2144.

818 Kessler A. & Halitschke R. (2007) Specificity and complexity: the impact of herbivore-  
819 induced plant responses on arthropod community structure. *Current Opinion in*  
820 *Plant Biology*, **10**, 409-414.

821 Kessler A. & Halitschke R. (2009) Testing the potential for conflicting selection on floral  
822 chemical traits by pollinators and herbivores: predictions and case study.  
823 *Functional Ecology*, **23**, 901-912.

824 Kessler A., Halitschke R. & Poveda K. (2011) Herbivory-mediated pollinator limitation:  
825 Negative impacts of induced volatiles on plant-pollinator interactions. *Ecology*,  
826 **92**, 1769-1780.

827 Kessler D., Diezel C. & Baldwin I.T. (2010) Changing pollinators as a means of escaping  
828 herbivores. *Current Biology*, **20**, 237-242.

829 Lehtilä K. & Strauss S.Y. (1997) Leaf damage by herbivores affects attractiveness to  
830 pollinators in wild radish, *Raphanus raphanistrum*. *Oecologia*, **111**, 396-403.

831 Leimu R., Muola A., Laukkanen L., Kalske A., Prill N. & Mutikainen P. (2012)  
832 Plant-herbivore coevolution in a changing world. *Entomologia Experimentalis et*  
833 *Applicata*, **144**, 3-13.

834 Lohman D., Zangerl A. & Berenbaum M. (1996) Impact of floral herbivory by parsnip  
835 webworm (Oecophoridae: *Depressaria pastinacella* Duponchel) on pollination  
836 and fitness of wild parsnip (Apiaceae: *Pastinaca sativa* L.). *American Midland*  
837 *Naturalist*, 407-412.

838 Lucas-Barbosa D., van Loon J.J.A. & Dicke M. (2011) The effects of herbivore-induced  
 839 plant volatiles on interactions between plants and flower-visiting insects.  
 840 *Phytochemistry*, **72**, 1647-1654.  
 841 Lüring M. & Scheffer M. (2007) Info-disruption: pollution and the transfer of chemical  
 842 information between organisms. *Trends in Ecology & Evolution*, **22**, 374-379.  
 843 McCall A.C. & Irwin R.E. (2006) Florivory: the intersection of pollination and herbivory.  
 844 *Ecology Letters*, **9**, 1351-1365.  
 845 Meiners T., Hacker N., Anderson P. & Hilker M. (2005) Response of the elm leaf beetle to  
 846 host plants induced by oviposition and feeding: the infestation rate matters.  
 847 *Entomologia Experimentalis et Applicata*, **115**, 171-177.  
 848 Mithöfer A. & Boland W. (2008) Recognition of herbivory-associated molecular patterns.  
 849 *Plant Physiology*, **146**, 825-831.  
 850 Mithöfer A. & Boland W. (2012) Plant defense against herbivores: chemical aspects.  
 851 *Annual Review of Plant Biology*, **63**, 431-450.  
 852 Mithöfer A., Wanner G. & Boland W. (2005) Effects of feeding *Spodoptera littoralis* on  
 853 lima bean leaves. II. Continuous mechanical wounding resembling insect feeding  
 854 is sufficient to elicit herbivory-related volatile emission. *Plant Physiology*, **137**,  
 855 1160-1168.  
 856 Mori N. & Yoshinaga N. (2011) Function and evolutionary diversity of fatty acid amino  
 857 acid conjugates in insects. *Journal of Plant Interactions*, **6**, 103-107.  
 858 Mumm R. & Dicke M. (2010) Variation in natural plant products and the attraction of  
 859 bodyguards involved in indirect plant defense. *Canadian Journal of Zoology*, **88**,  
 860 628-667.

Nottingham S.F., Hardie J., Dawson G.W., Hick A.J., Pickett J.A., Wadhams L.J. & Woodcock  
 C.M. (1991) Behavioral and electrophysiological responses of aphids to host and  
 nonhost plant volatiles. *Journal of Chemical Ecology*, **17**, 1231-1242.

Pare P.W., Farag M.A., Krishnamachari V., Zhang H., Ryu C.-M. & Kloepper J.W. (2005)  
 Elicitors and priming agents initiate plant defense responses. *Photosynthesis  
 Research*, **85**, 149-159.

Pearse I.S., Harris D.J., Karban R. & Sih A. (2013) Predicting novel herbivore–plant  
 interactions. *Oikos*, **122**, 1554-1564.

Pearson D.E. & Callaway R.M. (2005) Indirect nontarget effects of host-specific biological  
 control agents: Implications for biological control. *Biological Control*, **35**, 288-  
 298.

Peñuelas J. & Llusà J. (2004) Plant VOC emissions: making use of the unavoidable.  
*Trends in Ecology & Evolution*, **19**, 402-404.

Pichersky E. & Gershenzon J. (2002) The formation and function of plant volatiles:  
 perfumes for pollinator attraction and defense. *Current Opinion in Plant Biology*,  
**5**, 237-243.

Pierre P.S., Jansen J.J., Hordijk C.A., van Dam N.M., Cortesero A.-M. & Dugravot S. (2011)  
 Differences in volatile profiles of turnip plants subjected to single and dual  
 herbivory above-and belowground. *Journal of Chemical Ecology*, **37**, 368-377.

Poelman E.H., Broekgaarden C., Van Loon J.J.A. & Dicke M. (2008) Early season herbivore  
 differentially affects plant defence responses to subsequently colonizing  
 herbivores and their abundance in the field. *Molecular Ecology*, **17**, 3352-3365.

Poelman E.H., Bruinsma M., Zhu F., Weldegergis B.T., Boursault A.E., Jongema Y., van  
 Loon J.J., Vet L.E., Harvey J.A. & Dicke M. (2012) Hyperparasitoids use herbivore-

885 induced plant volatiles to locate their parasitoid host. *PLoS Biology*, **10**,  
 886 e1001435.  
 887 Poelman E.H., Zheng S.-J., Zhang Z., Heemskerk N.M., Cortesero A.-M. & Dicke M. (2011)  
 888 Parasitoid-specific induction of plant responses to parasitized herbivores affects  
 889 colonization by subsequent herbivores. *Proceedings of the National Academy of*  
 890 *Sciences*, **108**, 19647-19652.  
 891 Poveda K., Steffan-Dewenter I., Scheu S. & Tschardt T. (2003) Effects of below-and  
 892 above-ground herbivores on plant growth, flower visitation and seed set.  
 893 *Oecologia*, **135**, 601-605.  
 894 Quicke D.L. (1997) *Parasitic wasps*. Chapman & Hall Ltd.  
 895 Radville L., Chaves A. & Preisser E.L. (2011) Variation in plant defense against invasive  
 896 herbivores: evidence for a hypersensitive response in eastern hemlocks (*Tsuga*  
 897 *canadensis*). *Journal of Chemical Ecology*, **37**, 592-597.  
 898 Raffa K.F., Phillips T.W. & Salom S.M. (1993) Strategies and mechanisms of host  
 899 colonization by bark beetles. In: *Beetle-pathogen interactions in conifer forests*  
 900 (eds T.O. Schowalter & G. Filip), pp. 103-128. Academic Press, New York.  
 901 Raguso R.A. (2008) Wake up and smell the roses: the ecology and evolution of floral  
 902 scent. *Annual Review of Ecology, Evolution, and Systematics*, **39**, 549-569.  
 903 Rasman S., Köllner T.G., Degenhardt J., Hiltbold I., Toepfer S., Kuhlmann U., Gershenzon  
 904 J. & Turlings T.C. (2005) Recruitment of entomopathogenic nematodes by insect-  
 905 damaged maize roots. *Nature*, **434**, 732-737.  
 906 Reaser J.K., Meyerson L.A., Cronk Q., De Poorter M., Eldrege L., Green E., Kairo M., Latasi  
 907 P., Mack R.N. & Mauremootoo J. (2007) Ecological and socioeconomic impacts of  
 908 invasive alien species in island ecosystems. *Environmental Conservation*, **34**, 98-  
 909 111.

910 Reymond P. (2013) Perception, signaling and molecular basis of oviposition-mediated  
 911 plant responses. *Planta*, **238**, 247-258.

912 Riffell J.A. (2011) The neuroecology of a pollinator's buffet: olfactory preferences and  
 913 learning in insect pollinators. *Integrative and Comparative Biology*, **51**, 781-793.

914 Robert C.A.M., Erb M., Hibbard B.E., French B.W., Zwahlen C. & Turlings T.C.J. (2012) A  
 915 specialist root herbivore reduces plant resistance and uses an induced plant  
 916 volatile to aggregate in a density-dependent manner. *Functional Ecology*, **26**,  
 917 1429-1440.

918 Sakai A.K., Allendorf F.W., Holt J.S., Lodge D.M., Molofsky J., With K.A., Baughman S., Cabin  
 919 R.J., Cohen J.E. & Ellstrand N.C. (2001) The population biology of invasive species.  
 920 *Annual Review of Ecology and Systematics*, 305-332.

921 Schiestl F.P. (2010) The evolution of floral scent and insect chemical communication.  
 922 *Ecology Letters*, **13**, 643-656.

923 Schiestl F.P. & Johnson S.D. (2013) Pollinator-mediated evolution of floral signals. *Trends*  
 924 *in Ecology & Evolution*.

925 Shiojiri K., Takabayashi J., Yano S. & Takafuji A. (2002) Oviposition preferences of  
 926 herbivores are affected by tritrophic interaction webs. *Ecology Letters*, **5**, 186-  
 927 192.

928 Smallegange R., Van Loon J., Blatt S., Harvey J., Agerbirk N. & Dicke M. (2007) Flower vs.  
 929 leaf feeding by *Pieris brassicae*: glucosinolate-rich flower tissues are preferred  
 930 and sustain higher growth rate. *Journal of Chemical Ecology*, **33**, 1831-1844.

931 Snoeren T.A., Mumm R., Poelman E.H., Yang Y., Pichersky E. & Dicke M. (2010) The  
 932 herbivore-induced plant volatile methyl salicylate negatively affects attraction of  
 933 the parasitoid *Diadegma semiclausum*. *Journal of Chemical Ecology*, **36**, 479-489.

934 Sorte C.J., Williams S.L. & Zerebecki R.A. (2010) Ocean warming increases threat of  
 935 invasive species in a marine fouling community. *Ecology*, **91**, 2198-2204.  
 936 Tamiru A., Bruce T.J., Woodcock C.M., Caulfield J.C., Midega C.A., Ogot C.K., Mayon P.,  
 937 Birkett M.A., Pickett J.A. & Khan Z.R. (2011) Maize landraces recruit egg and  
 938 larval parasitoids in response to egg deposition by a herbivore. *Ecology Letters*,  
 939 **14**, 1075-1083.  
 940 Ton J., D'Alessandro M., Jourdie V., Jakab G., Karlen D., Held M., Mauch-Mani B. & Turlings  
 941 T.C. (2007) Priming by airborne signals boosts direct and indirect resistance in  
 942 maize. *The Plant Journal*, **49**, 16-26.  
 943 Tumlinson J.H. & Engelberth J. (2008) Fatty acid-derived signals that induce or regulate  
 944 plant defenses against herbivory. In: *Induced Plant Resistance to Herbivory*, pp.  
 945 389-407. Springer.  
 946 Turlings T.C., Tumlinson J.H. & Lewis W. (1990) Exploitation of herbivore-induced plant  
 947 odors by host-seeking parasitic wasps. *Science*, **250**, 1251-1253.  
 948 Turlings T.C., Wäckers F.L., Vet L.E., Lewis W.J. & Tumlinson J.H. (1993) Learning of host-  
 949 finding cues by hymenopterous parasitoids. In: *Insect learning*, pp. 51-78.  
 950 Springer.  
 951 Vet L.E. & Dicke M. (1992) Ecology of infochemical use by natural enemies in a tritrophic  
 952 context. *Annual Review of Entomology*, **37**, 141-172.  
 953 Vet L.E., Lewis W.J. & Carde R.T. (1995) Parasitoid foraging and learning. In: *Chemical*  
 954 *Ecology of Insects 2* (eds R.T. Cardé & W.J. Bell), pp. 65-101. Springer.  
 955 Vilà M., Espinar J.L., Hejda M., Hulme P.E., Jarošík V., Maron J.L., Pergl J., Schaffner U., Sun  
 956 Y. & Pyšek P. (2011) Ecological impacts of invasive alien plants: a meta-analysis  
 957 of their effects on species, communities and ecosystems. *Ecology Letters*, **14**, 702-  
 958 708.



959 Visser J. & Avé D. (1978) General green leaf volatiles in the olfactory orientation of the  
 960 Colorado beetle, *Leptinotarsa decemlineata*. *Entomologia Experimentalis et*  
 961 *Applicata*, **24**, 738-749.  
 962 Vos M., Berrocal S.M., Karamaouna F., Hemerik L. & Vet L. (2001) Plant-mediated  
 963 indirect effects and the persistence of parasitoid-herbivore communities. *Ecology*  
 964 *Letters*, **4**, 38-45.  
 965 Walther G.-R., Roques A., Hulme P.E., Sykes M.T., Pyšek P., Kühn I., Zobel M., Bacher S.,  
 966 Botta-Dukát Z. & Bugmann H. (2009) Alien species in a warmer world: risks and  
 967 opportunities. *Trends in Ecology & Evolution*, **24**, 686-693.  
 968 Webster B., Qvarfordt E., Olsson U. & Glinwood R. (2013) Different roles for innate and  
 969 learnt behavioral responses to odors in insect host location. *Behavioral Ecology*,  
 970 **24**, 366-372.  
 971 White E.M., Wilson J.C. & Clarke A.R. (2006) Biotic indirect effects: a neglected concept in  
 972 invasion biology. *Diversity and Distributions*, **12**, 443-455.  
 973 Winter T.R. & Rostás M. (2010) Nitrogen deficiency affects bottom-up cascade without  
 974 disrupting indirect plant defense. *Journal of Chemical Ecology*, **36**, 642-651.  
 975 Woodard A.M., Ervin G.N. & Marsico T.D. (2012) Host plant defense signaling in response  
 976 to a coevolved herbivore combats introduced herbivore attack. *Ecology and*  
 977 *Evolution*, **2**, 1056-1064.  
 978 Wright G.A. & Schiestl F.P. (2009) The evolution of floral scent: the influence of olfactory  
 979 learning by insect pollinators on the honest signalling of floral rewards.  
 980 *Functional Ecology*, **23**, 841-851.  
 981 Wu J. & Baldwin I.T. (2010) New insights into plant responses to the attack from insect  
 982 herbivores. *Annual Review of Genetics*, **44**, 1-24.

983 Yoneya K., Kugimiya S. & Takabayashi J. (2009) Can herbivore-induced plant volatiles  
984 inform predatory insect about the most suitable stage of its prey? *Physiological*  
985 *Entomology*, **34**, 379-386.

986 Yoshinaga N., Alborn H.T., Nakanishi T., Suckling D.M., Nishida R., Tumlinson J.H. & Mori  
987 N. (2010) Fatty acid-amino acid conjugates diversification in lepidopteran  
988 caterpillars. *Journal of Chemical Ecology*, **36**, 319-325.

## Tables


**Table 1.** Examples of interferences in infochemical networks.

Type of interaction	Organisms involved	Origin of the interference	Result of the interference	Reference
Herbivore-herbivore and Predator-prey	Parasitic wasp-caterpillars	Native herbivore species	Presence of a non-host herbivore decreases the attractiveness of host-infested plants for the parasitic wasp. Host herbivores favour plant infested by the non host for oviposition.	Shiojiri et al. 2002
Predator-prey	Zooplankton - alga	Pollutant	Presence of a surfactant induces a costly anti-predator aggregation behaviour in a green alga	Lürling 2007
Predator-prey	Predatory mite – Spider mites	Uncommon herbivore species	Presence of the uncommon herbivore decreases the attractiveness of plants damaged by suitable prey	Fadini et al. 2010
Predator-prey	Parasitic wasp-caterpillar	Native herbivore species	Presence of a non-host herbivore increases the attractiveness of host-infested plants, but decreases the searching efficiency of the parasitoid.	Bukovinszky et al. 2012
Pollinator-flower	Wild tomato plants-bees	Native herbivore species	Plants damaged by herbivores show reduced attractiveness to pollinators and suffer reduced seed sets.	Kessler et al. 2011
Pollinator-flower	Wild radish-syrphid flies	Native herbivore species	Lower pollinator visitation rates on plants damaged by herbivores	Lehtilä and Strauss 1997


998  
999 **Figure legends**

1000 **Figure 1.** Illustration of how invasive insect herbivore may disrupt infochemical  
1001 networks through interferences in herbivore induced plant volatiles (illustration by  
1002 Thomas Degen).


1003 **Figure 2.** Illustration of conceivable plant responses to exotic herbivory. Letters  
1004 represent volatile compounds emitted in response to mechanical damage or to feeding  
1005 by two native herbivores. Three types of plant response to exotic herbivory are  
1006 proposed. **Null response:** the blend emitted resembles the blend emitted after  
1007 mechanical damage. **Identical response:** the blend emitted resembles the blend emitted  
1008 in response to one native species (here, herbivore #1). **Novel response:** the blend  
1009 emitted diverges from the blend emitted in response to native herbivores.



hypothetical BVOC infochemical web after invasion



native interaction web and BVOC infochemical web



hypothetical BVOC infochemical web after invasion



